

A RAMSEY SPACE OF INFINITE POLYHEDRA AND THE INFINITE RANDOM POLYHEDRON

JOSÉ G. MIJARES AND GABRIEL PADILLA

ABSTRACT. In this paper we introduce a new topological Ramsey space (see [19]) whose elements are infinite ordered polyhedra. Then, we show as an application that the set of finite polyhedra satisfies two types of Ramsey property: one in the sense of Graham, Leeb & Rothschild in [5], when viewed as a category over $\mathbb{N}$; the other in the sense of [8, 12, 16] (for instance), when considered as a class of finite structures. Finally, it is proven that the group of automorphism of the Fraïssé limit of the class of finite ordered polyhedra is extremely amenable. We call this Fraïssé limit the *infinite (ordered) random polyhedron*.

INTRODUCTION

A polyhedron is a geometric object built up through a finite number of suitable amalgamations of convex hulls of finite sets; polyhedrons are generated in this way by simplexes. Simplicial morphisms are locally linear maps that preserve vertices. An ordered polyhedron is a polyhedron for which we have imposed a linear order on the set of its vertices. As we only consider order-preserving morphisms, ordered polyhedra are rigid, i. e., admit no non-trivial automorphisms; this is an easy consequence of the well order principle. In this paper we define a new topological Ramsey space (see [19]) whose elements are essentially infinite ordered polyhedra.

The prototypical topological Ramsey space is due to Ellentuck [4]. It is the set $\mathbb{N}^{[\infty]}$ of infinite subsets of $\mathbb{N}$, equipped with the *exponential topology*, whose basic sets are of the form:

$$(1) \quad [a, A] = \{B \in \mathbb{N}^{[\infty]} : a \sqsubset B \text{ \& } B \subseteq A\}.$$

where a is a finite subset of $\mathbb{N}$ and $A \in \mathbb{N}^{[\infty]}$. Here, $a \sqsubset B$ means that a is an initial segment of B . Recall that a set $\mathcal{X} \subseteq \mathbb{N}^{[\infty]}$ is *Ramsey* if for every neighborhood $[a, A] \neq \emptyset$ there exists $B \in [a, A]$ such that $[a, B] \subseteq \mathcal{X}$ or $[a, B] \cap \mathcal{X} = \emptyset$. And it is *Ramsey null* if for every neighborhood $[a, A]$ there exists $B \in [a, A]$ such that $[a, B] \cap \mathcal{X} = \emptyset$. The main result in [4] states the following:

Theorem 0.1. (*Ellentuck [4], 1974*) *A set $\mathcal{X} \subseteq \mathbb{N}^{[\infty]}$ is Ramsey if and only if it has the Baire property with respect to the exponential topology. And $\mathcal{X}$ is Ramsey null if and only if it is meager with respect to the exponential topology.*

The theory of topological Ramsey spaces is developed in [3, 19] following the work of Ellentuck's. In the following section we will present the fundamental concepts

Date: 07/07/2012.

1991 Mathematics Subject Classification. 05D10, 52B05.

Key words and phrases. Ramsey spaces, random polyhedra.

of that theory. In Section 2 we will define our new topological Ramsey space $\mathcal{P}$ and in the rest of the paper we will use $\mathcal{P}$ to study the Ramsey property for the category of finite ordered polyhedra and prove that the group of automorphism of the Fraïssé limit of the class of finite ordered polyhedra is extremely amenable. Finally, a description of this limit is given. We call it the *infinite (ordered) random polyhedron*.

1. RAMSEY SPACES

The definitions and results throughout this section are taken from [19]. A previous presentation can also be found in [3]. Consider a triplet of the form $(\mathcal{R}, \leq, r)$, where $\mathcal{R}$ is a set, $\leq$ is a quasi order on $\mathcal{R}$ and $r : \mathbb{N} \times \mathcal{R} \rightarrow \mathcal{AR}$ is a function with range $\mathcal{AR}$. For every $n \in \mathbb{N}$ and every $A \in \mathcal{R}$, let us write $r_n(A) := r(n, A)$ and $\mathcal{AR}_n := \{r_n(A) : A \in \mathcal{R}\}$. We say that $r_n(A)$ is **the n th approximation of A** . In order to capture the combinatorial structure required to ensure the provability of an Ellentuck type Theorem, some assumptions on $(\mathcal{R}, \leq, r)$ will be imposed. The first is the following:

(A.1)

(A.1.1) For any $A \in \mathcal{R}$, $r_0(A) = \emptyset$.

(A.1.2) For any $A, B \in \mathcal{R}$, if $A \neq B$ then $(\exists n) (r_n(A) \neq r_n(B))$.

(A.1.3) If $r_n(A) = r_m(B)$ then $n = m$ and $(\forall i < n) (r_i(A) = r_i(B))$.

A.1 allows us to identify each $A \in \mathcal{R}$ with the sequence $(r_n(A))_n$ of its approximations. In this way, if $\mathcal{AR}$ has the discrete topology, $\mathcal{R}$ can be identified with a subspace of the (metric) space $\mathcal{AR}^{\mathbb{N}}$ (with the product topology) of all the sequences of elements of $\mathcal{AR}$, see for instance [1, 7]. We will say that $\mathcal{R}$ is **metrically closed** if it is a closed subspace of $\mathcal{AR}^{\mathbb{N}}$. The basic open sets generating the metric topology on $\mathcal{R}$ inherited from the product topology of $\mathcal{AR}^{\mathbb{N}}$ are of the form:

$$(2) \quad [a] = \{B \in \mathcal{R} : (\exists n)(a = r_n(B))\}$$

where $a \in \mathcal{AR}$. For each $a \in \mathcal{AR}$ let us define the **length** of a , as the unique integer $|a| = n$ such that $a = r_n(A)$ for some $A \in \mathcal{R}$. The **Ellentuck type neighborhoods** are of the form:

$$(3) \quad [a, A] = \{B \in \mathcal{R} : (\exists n)(a = r_n(B)) \text{ and } (B \leq A)\}$$

where $a \in \mathcal{AR}$ and $A \in \mathcal{R}$. Let $\mathcal{AR}(A) = \{a \in \mathcal{AR} : [a, A] \neq \emptyset\}$. We also write $[n, A] := [r_n(A), A]$. Given a neighborhood $[a, A]$ and $n \geq |a|$, let $r_n[a, A]$ be the image of $[a, A]$ by the function r_n , i.e., the set $\{b \in \mathcal{AR} : \exists B \in [a, A] \text{ such that } b = r_n(B)\}$.

1.1. Ramsey sets. A set $\mathcal{X} \subseteq \mathcal{R}$ is **Ramsey** if for every neighborhood $[a, A] \neq \emptyset$ there exists $B \in [a, A]$ such that $[a, B] \subseteq \mathcal{X}$ or $[a, B] \cap \mathcal{X} = \emptyset$. A set $\mathcal{X} \subseteq \mathcal{R}$ is **Ramsey null** if for every neighborhood $[a, A]$ there exists $B \in [a, A]$ such that $[a, B] \cap \mathcal{X} = \emptyset$.

1.2. Topological Ramsey spaces. We say that $(\mathcal{R}, \leq, r)$ is a **topological Ramsey space** iff subsets of $\mathcal{R}$ with the Baire property are Ramsey and meager subsets of $\mathcal{R}$ are Ramsey null.

(A.2) [Finitization] There is a quasi order $\leq_{fin}$ on $\mathcal{AR}$ such that:

$$(i) \ A \leq B \text{ iff } (\forall n) (\exists m) (r_n(A) \leq_{fin} r_m(B)).$$

(ii) $\{b \in \mathcal{AR} : b \leq_{fin} a\}$ is finite, for every $a \in \mathcal{AR}$.

Given $A \in \mathcal{R}$ and $a \in \mathcal{AR}(A)$, we define the *depth of a in A* as

$$(4) \quad \text{depth}_A(a) := \min\{n : a \leq_{fin} r_n(A)\}.$$

Lemma 1.1. *Given $A \in \mathcal{R}$ and $a \in \mathcal{AR}(A)$, $|a| \leq \text{depth}_A(a)$.*

[Proof] By axioms A.1.3 and A.2.1. □

(A.3) [Amalgamation] Given a and A with $\text{depth}_A(a) = n$, the following holds:

- (i) $(\forall B \in [n, A]) \quad ([a, B] \neq \emptyset).$
- (ii) $(\forall B \in [a, A]) \quad (\exists A' \in [n, A]) \quad ([a, A'] \subseteq [a, B]).$

(A.4) [Pigeonhole Principle] Given a and A with $\text{depth}_A(a) = n$, for every $\mathcal{O} \subseteq \mathcal{AR}_{|a|+1}$ there is $B \in [n, A]$ such that $r_{|a|+1}[a, B] \subseteq \mathcal{O}$ or $r_{|a|+1}[a, B] \subseteq \mathcal{O}^c$.

Theorem 1.2 (See [3, 19]). **[Abstract Ellentuck Theorem]** *Any $(\mathcal{R}, \leq, r)$ with $\mathcal{R}$ metrically closed and satisfying (A.1)-(A.4) is a topological Ramsey space.*

Notation: The following is taken from [11]. For $k, m \in \mathbb{N}$, $A \in \mathcal{R}$ and $b \in \mathcal{AR}(A)$, let us define

$$(5) \quad \mathcal{AR}_k^m(A) := \{a \in \mathcal{AR}_k(A) : \text{depth}_A(a) = m\},$$

$$(6) \quad \mathcal{AR}_k^m(A, b) := \{a \in \mathcal{AR}_k^m(A) : a \leq_{fin} b\}$$

In [11], one can find a proof of the following abstract version of finite Ramsey's Theorem:

Theorem 1.3. *Let $A \in \mathcal{R}$ and $k, n, r \in \mathbb{N}$ be given. Then, there exists $m \in \mathbb{N}$ such that for every coloring $c : \mathcal{AR}_k^m(A) \rightarrow r$, there exists $b \in \mathcal{AR}_n^m(A)$ such that c is constant in $\mathcal{AR}_k^m(A, b)$.*

For instance, consider the triplet $\mathcal{E} = (\mathbb{N}^{[\infty]}, \subseteq, i)$, where $\subseteq$ is the inclusion relation and $i : \mathbb{N} \times \mathbb{N}^{[\infty]} \rightarrow \mathbb{N}^{[<\infty]}$, is the approximation function:

$$(7) \quad i(n, A) = \text{the first } n \text{ elements of } A$$

In this case, the instance of the abstract Ellentuck theorem is Theorem 0.1 and the instance of Theorem 1.3 if the finite version of Ramsey's theorem [18].

From now on, we will refer to $\mathcal{E} = (\mathbb{N}^{[\infty]}, \subseteq, i)$ as **Ellentuck's space**.

2. THE TOPOLOGICAL RAMSEY SPACE $\mathcal{P}$

In this Section we will consider pairs (x, S_x) satisfying the following:

- (1) $x \subseteq \mathbb{N}$,
- (2) $S_x \subseteq x^{[<\infty]}$ is **hereditary**, i.e., $u \subseteq v$ & $v \in S_x \Rightarrow u \in S_x$, and
- (3) $\bigcup S_x = \bigcup \{u : u \in S_x\} = x$.

Given a pair (x, S_x) satisfying the conditions above and $y \subseteq x$, we let $S_x \upharpoonright y = \{u \cap y : u \in S_x\}$. Given two pairs (x, S_x) , (y, S_y) satisfying the conditions (1), (2) and (3); we write

$$(8) \quad (y, S_y) \leq (x, S_x) \Leftrightarrow y \subseteq x \text{ \& } S_y \subseteq S_x.$$

For instance, if $y \subset x$ then $(y, S_x \upharpoonright y) \leq (x, S_x)$. In particular, if $n \in \mathbb{N}$ and $|x| \geq n$ let $x \upharpoonright_n$ be the set of the first n elements of x and $S_{x \upharpoonright_n} = S_x \upharpoonright (x \upharpoonright_n)$. The pair

$$(9) \quad r_n(x, S_x) = (x \upharpoonright_n, S_{x \upharpoonright_n})$$

is the n th **approximation** of (x, S_x) . Notice that $r_n((x, S_x)) \leq (x, S_x)$ for all $n \leq |x|$.

Write $\mathcal{AP}$ to denote the set of all pairs (x, S_x) such that $|x| < \infty$; and $\mathcal{P}$ to denote its complement, whose elements will be written $(A, S_A), (B, S_B) \dots$ in capital letters. There is a well defined surjective function

$$(10) \quad \mathcal{P} \times \mathbb{N} \xrightarrow{r} \mathcal{AP} \quad r((A, S_A), n) = r_n(A, S_A)$$

In the rest of this section we shall prove the following:

Theorem 2.1. $(\mathcal{P}, \leq, r)$ is a topological Ramsey space.

The proof of Theorem 2.1 will be divided into several lemmas, showing that $(\mathcal{P}, \leq, r)$ satisfies the conditions of the abstract Ellentuck theorem.

Lemma 2.2. $(\mathcal{P}, \leq, r)$ satisfies axiom **A.1**

- (1) For every $(A, S_A) \in \mathcal{P}$, $r_0((A, S_A)) = \emptyset$.
- (2) If $(A, S_A) \neq (B, S_B)$ then there exists n such that $r_n((A, S_A)) \neq r_n((B, S_B))$.
- (3) If $r_n((A, S_A)) = r_m((B, S_B))$ then $n = m$ and for every $i < n$, $r_i((A, S_A)) = r_i((B, S_B))$.

[Proof] Straightforward. □

Hence each element of $\mathcal{P}$ can be identified with the sequence of its approximations. Next we consider $\mathcal{P}$ as a subset of the product space $\mathcal{AP}^{\mathbb{N}}$, regarding $\mathcal{AP}$ as a discrete space.

Lemma 2.3. $\mathcal{P}$ is a closed subset of $\mathcal{AP}^{\mathbb{N}}$.

[Proof] By Lemma §2.2 the injection $\mathcal{P} \xrightarrow{\varphi} \mathcal{AP}^{\mathbb{N}}$ given by

$$\varphi(A, S_A) = (r_0(A, S_A), r_1(A, S_A), \dots)$$

is continuous. Let us show that $\varphi(\mathcal{P})$ is closed. For a closure point $\alpha = \{(a^j, S_{a^j})\}_{j \in \mathbb{N}}$ in $\overline{\varphi(\mathcal{P})} \subset \mathcal{AP}^{\mathbb{N}}$ and a sequence $\{(A^k, S_{A^k})\}_{k \in \mathbb{N}}$ in $\mathcal{P}$; if $\{\varphi(A^k, S_{A^k})\}_{k \in \mathbb{N}}$ converges to α then

$$(\forall n \in \mathbb{N}) (\exists k_n \in \mathbb{N}) \quad k \geq k_n \Rightarrow (\forall j \leq n) \quad r_j(A^k, S_{A^k}) = (a^j, S_{a^j})$$

Taking a strictly increasing sequence $k_n < k_{n+1} \forall n \in \mathbb{N}$ we get

$$n > m \Rightarrow r_m(A^{k_n}, S_{A^{k_n}}) = (a^{k_m}, S_{a^{k_m}})$$

Define $A = \bigcup_{n \in \mathbb{N}} a^{k_n}$ and $S_A = \bigcup_{n \in \mathbb{N}} S_{a^{k_n}}$. Then $(A, S_A) \in \mathcal{P}$ and $\varphi(A, S_A) = \alpha$ by construction. □

Let us define the preorder $\leq_{fin}$ and the partial order $\sqsubseteq$ on $\mathcal{AP}$ as follows:

$$(11) \quad (a, S_a) \leq_{fin} (b, S_b) \Leftrightarrow (a, S_a) \leq (b, S_b) \text{ \& \; } \max(a) = \max(b)$$

$$(12) \quad (a, S_a) \sqsubseteq (b, S_b) \Leftrightarrow a \sqsubseteq b \text{ \& \; } (a, S_a) \leq (b, S_b)$$

Here we are using the same symbol " $\sqsubseteq$ " to indicate that the set a is an initial segment of the set b . The following two lemmas are straightforward; we leave the details to the reader.

Lemma 2.4. $(\mathcal{P}, \leq, r)$ satisfies axiom **A.2**

- (1) If $(A, S_A) \leq (B, S_B)$ then $\forall n \exists m, r_n((A, S_A)) \leq_{fin} r_m((B, S_B))$.
- (2) For every $(a, S_a) \in \mathcal{AP}$ the set $\{(b, S_b) : (b, S_b) \leq_{fin} (a, S_a)\}$ is finite.
- (3) If $(a, S_a) \leq_{fin} (b, S_b)$ and $(c, S_c) \sqsubseteq (a, S_a)$ then there is $(d, S_d) \sqsubseteq (b, S_b)$ such that $(c, S_c) \leq_{fin} (d, S_d)$.

Lemma 2.5. $(\mathcal{P}, \leq, r)$ satisfies axiom **A.3**

Let $n = \text{depth}_{(B, S_B)}((a, S_a))$.

- (1) If $(A, S_A) \in [n; (B, S_B)]$ then $[(a, S_a); (A, S_A)] \neq \emptyset$.
- (2) For every $(A, S_A) \in [(a, S_a); (B, S_B)]$ there exists $(A', S_{A'}) \in [n; (B, S_B)]$ such that $\emptyset \neq [(a, S_a); (A', S_{A'})] \subseteq [(a, S_a); (A, S_A)]$.

For every natural number n , let

$$\mathcal{AP}_n := \{(a, S_a) \in \mathcal{AP} : |a| = n\}.$$

If $(a, S_a) \in \mathcal{AP}_n$, then we say that the **length** of (a, S_a) is n or simply write $|(a, S_a)| = n$.

Finally, we prove the following

Lemma 2.6. *Pigeonhole principle A.4 for $(\mathcal{P}, \leq, r)$:*

Let $n = \text{depth}_{(B, S_B)}((a, S_a))$, $k = |(a, S_a)|$ and $c : \mathcal{AP}_{k+1} \rightarrow \{0, 1\}$ be any partition. There exists $(A, S_A) \in [n; (B, S_B)]$ such that c is constant in $r_{k+1}[(a, S_a); (A, S_A)]$.

[Proof] Let

$$X = \{b \in B/a : \text{there exists } u \in S_a \text{ such that } u \cup \{b\} \in S_B\}$$

For $i \in \{0, 1\}$, let

$$X_i = \{b \in X : c((a \cup \{b\}, S_B \upharpoonright a \cup \{b\})) = i\}.$$

By the classical pigeon hole principle, there is $i_0 \in \{0, 1\}$ such that $|X_{i_0}| = \infty$. So let

$$A = (B \upharpoonright n) \cup X_{i_0} \text{ and } S_A = S_B \upharpoonright A$$

Then $(A, S_A) \in [n; (B, S_B)]$ is as required. $\square$

Now we can proof that $(\mathcal{P}, \leq, r)$ is a topological Ramsey space:

[Proof of Theorem 2.1] In virtue of the abstract Ellentuck theorem, the required result follows from Lemmas 2.2, 2.3, 2.4, 2.5, 2.6. $\square$

Remark 2.7. (Ellentuck's space as a subspace of $\mathcal{P}$) Notice that we can identify each $A \in \mathbb{N}^{[\infty]}$ with the pair $(A, \{\{x\} : x \in A\} \cup \{\emptyset\})$. In this way, we can view Ellentuck's space $\mathcal{E}$ as a (closed) subspace of $\mathcal{P}$.

Now we give an alternative proof to the well known fact that $\mathcal{E}$ is a Ramsey space. Recall the approximation function $i : \mathbb{N} \times \mathbb{N}^{[\infty]} \rightarrow \mathbb{N}^{[<\infty]}$, given by

$$i(n, A) = \text{the first } n \text{ elements of } A.$$

Corollary 2.8. (Ellentuck [4], 1974) $\mathcal{E} = (\mathbb{N}^{[\infty]}, \subseteq, i)$ is a topological Ramsey space.

[*Proof*] Fix $\mathcal{X} \subseteq \mathbb{N}^{[\infty]}$ with the Baire property with respect to the exponential topology of $\mathcal{E}$. Since $\mathcal{E}$ as a closed subspace of $\mathcal{P}$, it is easy to show that $\mathcal{X}$ has the Baire property with respect to the Ellentuck-like topology of $\mathcal{P}$ ⁽¹⁾. Given a nonempty neighborhood $[a, A]$ in $\mathcal{E}$, let $S_a = \{\{x\} : x \in a\} \cup \{\emptyset\}$ and $S_A = \{\{x\} : x \in A\} \cup \{\emptyset\}$. Then, consider the neighborhood $[(a, S_a), (A, S_A)]$ in $\mathcal{P}$. Applying Theorem 2.1 we obtain $(B, S_B) \in [(a, S_a), (A, S_A)]$ such that $[(a, S_a), (B, S_B)] \subseteq \mathcal{X}'$ or $[(a, S_a), (B, S_B)] \cap \mathcal{X}' = \emptyset$, where

$$\mathcal{X}' = \{(A, \{\{x\} : x \in A\} \cup \{\emptyset\}) : A \in \mathcal{X}\}.$$

Notice that, by necessity, $S_B = \{\{x\} : x \in B\} \cup \{\emptyset\}$. Hence, $[a, B] \subseteq \mathcal{X}$ or $[a, B] \cap \mathcal{X} = \emptyset$.

If $\mathcal{X}$ is meager with respect to the exponential topology of $\mathcal{E}$ then the same argument works but in addition the case $[a, B] \subseteq \mathcal{X}$ will never happen, by the meagerness of $\mathcal{X}$. This completes the proof. $\square$

We will see later in Theorem 4.1 that we can understand $\mathcal{AP}$ as a category (over $\mathbb{N}$) with suitable embeddings, and that it is in fact a **Ramsey category** in the sense of [5]. We will also see that it is a **Ramsey class of finite structures** in the sense of [8, 12, 16] (for instance). But before that, we will give a geometric interpretation of the Ramsey space $(\mathcal{P}, \leq, r)$.

3. THE RAMSEY SPACE $\mathcal{P}$ GEOMETRICALLY INTERPRETED

3.1. Simplexes. Let $V = \{v_0, \dots, v_n\} \subset \mathbb{R}^d$ be an affinely independent finite subset of some euclidean space, which we call the **ambient space**. The n -**simplex** generated by V , written $\langle V \rangle$, is the convex hull generated by the points of V ; i.e. the set of convex combinations,

$$\sum_{i=0}^n a_i v_i \quad a_0, \dots, a_n \in [0, \infty); \quad a_0 + \dots + a_n = 1$$

A **vertex** of $\langle V \rangle$ is a point in V . The integer n is the affine **dimension** of $\langle V \rangle$. Thus

$\dim(\langle V \rangle) + 1 = |V|$. A linear **morphism** between simplexes $\langle W \rangle \xrightarrow{f} \langle V \rangle$ is the restriction of a linear function f between the respective ambient spaces, such that $f(\langle W \rangle) \subset \langle V \rangle$. An **embedding** is a vertex-preserving linear injective morphism, i.e. $f(W) \subset V$. For $0 \leq m \leq n$, a m -**face** (or m -**subsimplex**) of $\langle V \rangle$ is the image

$\rho(f) \subset \langle V \rangle$ of an m -simplex $\langle W \rangle \xrightarrow{f} \langle V \rangle$ through an embedding $\langle W \rangle \xrightarrow{f} \langle V \rangle$. Notice that the faces of V can be obtained from the subsets of V . By convention we will consider $\emptyset$ as the empty face with affine dimension 0. Given $n \in \mathbb{N}$, we write

$$\Delta^n = \langle \{e_0, \dots, e_n\} \rangle \subset \mathbb{R}^{n+1}$$

for the **standard n -simplex** generated by the canonical basis.

¹ This can be deduced from two facts: (a) Since $\mathcal{E}$ is closed in $\mathcal{P}$, every meager subset of $\mathcal{E}$ is still meager in $\mathcal{P}$; and (b) Subsets of $\mathcal{P}$ with the Baire property constitute a σ -algebra.

3.2. Simplicial sets and rigid polyhedra. A geometric polyhedron is the amalgamated sum of a countable collection of standard simplexes through a countable family of embeddings; a simplicial set is the prescription of these embeddings, so it is essentially a recipe for gluing simplexes. Simplicial sets always are induced by the category of finite ordinals; see [9, pp.171-174]. Here there is another approach: A (finite) **simplicial set** is a set $S \subset \mathbb{N}^{[<\infty]}$ such that $S = S_x$ for some $(x, S_x) \in \mathcal{AP}$. The elements of $x = \cup S$ correspond to the vertices in a geometric realization of S . For instance, if $|\cup S| < \infty$ define

$$(13) \quad V_S = \{e_j : j \in \cup S\} \subset \mathbb{R}^{\max(\cup S)+1}$$

as the corresponding subset of elements in the canonical basis of $\mathbb{R}^{\max(\cup S)+1}$. We generate from S a polyhedron K by taking the amalgamated sum of the family of simplexes $\{\langle V_u \rangle : u \text{ is maximal in } (S, \subset)\}$ glued by the (linear) embeddings

$$(14) \quad \langle V_u \rangle \longleftarrow \langle V_{u \cap v} \rangle \longrightarrow \langle V_v \rangle$$

induced by the inclusions $V_{u \cap v} \subset V_u$ and $V_{u \cap v} \subset V_v$. Notice that the geometric object K does not depend on V_S but rather on $(S, \subset)$. As an example, one could have decided to embed K in $\mathbb{R}^{|\cup S|+1}$ instead of $\mathbb{R}^{\max(\cup S)+1}$; this can be done by

taking any bijection $\cup S \xrightarrow{\sim} |\cup S|$, this leads to a polyhedron which is isomorphic to K . For instance, if $S_x = 2^x$ is the set of parts of x then $K = \Delta^{|\cup S|} \subset \mathbb{R}^{|\cup S|+1}$ is a standard simplex.

Conversely, given a geometric polyhedron K we get a simplicial set by letting $x = V_K$ be the set of vertices in K and

$$S_x = \{V \subset V_K : \langle V \rangle \subset K\}$$

In other words, $V \in S_x$ iff $\langle V \rangle$ is a face of K . Then $(x, S_x) \in \mathcal{AP}$ and $S = S_x$ is a simplicial set. From the above discussion, it should now be clear that each simplicial set $S = S_x$ coming from some pair $(x, S_x) \in \cup \mathcal{AP}$ represents, geometrically, the set of faces of some polyhedron K with finite dimension.

A (finite) **ordered polyhedron** is a geometric polyhedron for which we have prefixed a linear order on the set of its vertices. It corresponds to a pair $(x, S_x) \in \mathcal{AP}$.

An **embedding** $(x, S_x) \xrightarrow{f} (y, S_y)$ is an injective function $x \xrightarrow{f} y$ such that

$u \in S_x \Rightarrow f(u) \in S_y$. A **rigid embedding** is an embedding $(x, S_x) \xrightarrow{f} (y, S_y)$ such that f is order-preserving: $i < j \Rightarrow f(i) < f(j)$.

The following will be useful in the sequel:

Lemma 3.1. *Each finite polyhedron can be rigidly embedded in some simplex.*

[Proof] If $(x, S_x) \in \mathcal{AP}$ is a finite polyhedron let $n = \max(x)$ be the maximum of x . Then the inclusions $x \subset n$, $S_x \subset 2^n$ induce a rigid embedding

$$(x, S_x) \longrightarrow (n, 2^n)$$

Geometrically, this is just the inclusion of the polyhedron K determined by (x, S_x) in the standard simplex $\Delta^n \subset \mathbb{R}^{n+1}$. $\square$

3.3. The Ramsey space $\mathcal{P}$ geometrically interpreted. Let $\{\mathbf{e}_n : n \in \mathbb{N}\}$ be the canonical basis of the real separable Hilbert space l^2 . By the correspondence

$$(15) \quad u \mapsto V_u = \{\mathbf{e}_j : j \in u\} \quad \forall u \in S_A$$

each $(A, S_A) \in \mathcal{P}$ can be considered as an infinite-dimensional polyhedron $K \subset l^2$ for which we have fixed a linear order on its vertices. We call $\mathcal{P}$ **the Ramsey space of infinite countable ordered polyhedra**. There is also a notion of rigid embeddings in this case.

4. CATEGORICAL RAMSEY PROPERTIES FOR POLYHEDRA

4.1. Ramsey Categories. In this § we follow [5]. We say that $\mathfrak{D}$ is **pre-Ramsey** iff it satisfies the following properties.

- (1) $\text{Obj}(\mathfrak{D}) = \mathbb{N}$ is the set of non-negative integers.
- (2) For each pair of non-negative integers $m \leq n$ there is an integer $r_{m,n}$ such that $\binom{m}{n}_{\mathfrak{D}}$ is a finite set with $r_{m,n}$ elements. In particular, $r_{0,0} = 1$.
- (3) $\text{Mor}(\mathfrak{D}) \subset \mathcal{E}$, all morphisms of $\mathfrak{D}$ are embeddings ⁽²⁾.

A pre-Ramsey subcategory $\mathfrak{D}$ is **Ramsey** iff additionally it satisfies

- (4) The Ramsey property: Given $m, n, r \in \mathbb{N}$ there is $N_{(m,n,r)} \in \mathbb{N}$ such that for all $N \geq N_{(m,n,r)}$; every r -coloring $\binom{N}{m}_{\mathfrak{D}} \longrightarrow r$ has a monochromatic sub-object in $\text{Obj}(\mathfrak{D})$ of n ; ⁽³⁾.

4.2. Finite polyhedra as Ramsey Category. In virtue of Theorem 2.1, the following instance of Theorem 1.3 holds. This is the version of finite Ramsey's Theorem corresponding to the Ramsey space $\mathcal{P}$ (Recall the notation at the end of Section 1.2):

Theorem 4.1. *Let $(A, S_A) \in \mathcal{P}$ and $k, n, r \in \mathbb{N}$ be given. Then, there exists $m \in \mathbb{N}$ such that for every coloring $c : \mathcal{AP}_k^m((A, S_A)) \rightarrow r$, there exists $(b, S_b) \in \mathcal{AP}((A, S_A))^m_n$ such that c is constant in $\mathcal{AP}_k^m((A, S_A), (b, S_b))$.*

Thanks to Theorem 4.1, it is now easy to prove that the category of finite polyhedra is Ramsey. Notice that we do not need to check the conditions of sufficiency stated in the main Theorem [5, p.418] but simply translate Theorem 4.1 into its geometric/categorical version.

Theorem 4.2. [Categorical Ramsey property for polyhedra] *The category $\mathfrak{P}$ of finite ordered polyhedra and rigid embeddings is Ramsey.*

[Proof] Fix $k, n, r \in \mathbb{N}$ and apply Theorem 4.1 to $A = \mathbb{N}$ and $S_A = \mathbb{N}^{[<\infty]}$. $\square$

5. RAMSEY CLASSES

In this section we present some basic concepts on Ramsey classes, Fraïssé theory and extremely amenability of automorphism groups. For more details and indepth treatment of these subjects see [2, 8, 10, 16].

²For finite sets this means that all morphisms of $\mathfrak{D}$ are injective functions.

³We will assume the usual identification between each non-negative number r and the cardinal r . So $0 = \emptyset$ and $r = \{0, \dots, r-1\}$ are sets.

5.1. Structures. A **signature** $L = \langle (R_i)_{i \in I}, (F_j)_{j \in J} \rangle$ is a couple of two countable sets, we say that $(R_i)_{i \in I}$ is a set **relation symbols** while $(F_j)_{j \in J}$ is a set of **function symbols**. Each relation (resp. function) symbol R_i (resp. F_j) has an associated integer number $n(i) > 0$ (resp. $m(j) \geq 0$) called its **arity**.

Each signature L has an associated countable first order language which contains the symbols of L , a countable set of variable symbols (say $x_1, x_2, \dots$ etc.); the usual connective symbols ($\wedge, \vee, \neg, \rightarrow$), quantifiers ($\forall, \exists$) and a binary symbol for the equality ($=$). With little abuse of notation we will still refer to this language with the letter L ; it might also have constant symbols (function symbols with arity 0) though we will not insist on them.

Given a language L , a L -**structure**, $\mathbb{A} = \langle A, (R_i^{\mathbb{A}})_{i \in I}, (F_j^{\mathbb{A}})_{j \in J} \rangle$ is a triple constituted as follows:

- A non empty set $A \neq \emptyset$ called the **universe** of the structure;
- a set of relations $(R_i^{\mathbb{A}})_{i \in I}$ where $R_i^{\mathbb{A}} \subseteq A^{n(i)}$ for each $i \in I$; and
- a set of functions $(F_j^{\mathbb{A}})_{j \in J}$ where $F_j^{\mathbb{A}} : A^{m(j)} \longrightarrow A$ for each $j \in J$.

We say that $\mathbb{A}$ is a **relational structure** if L has no function symbols.

A **morphism** of L -structures $\mathbb{A} \xrightarrow{\pi} \mathbb{B}$ is a map $A \xrightarrow{\pi} B$ between the respective universe sets, such that for each relation symbol R_i and each function symbol F_j ; it satisfies

- $(a_1, \dots, a_{n(i)}) \in R_i^{\mathbb{A}}$ iff $(\pi(a_1), \dots, \pi(a_{n(i)})) \in R_i^{\mathbb{B}}$; for all $a_1, \dots, a_{n(i)} \in A$.
- $\pi(F_j^{\mathbb{A}}(a_1, \dots, a_{m(j)})) = F_j^{\mathbb{B}}(\pi(a_1), \dots, \pi(a_{m(j)}))$ for all $a_1, \dots, a_{m(j)} \in A$.

When π is bijective we say that it is an **isomorphism** and $\mathbb{A}, \mathbb{B}$ are **isomorphic** structures, or just $\mathbb{A} \cong \mathbb{B}$. When π is injective we say that it is an **embedding**. In particular, we say that $\mathbb{A}$ is a **substructure** of $\mathbb{B}$, and write $\mathbb{A} \leq \mathbb{B}$ whenever

$A \subseteq B$ and the inclusion map $\mathbb{A} \xrightarrow{i} \mathbb{B}$ is an embedding. In that case

- $R_i^{\mathbb{A}} = R_i^{\mathbb{B}} \cap A^{n(i)}$ for each $i \in I$.
- $F_j^{\mathbb{A}} = F_j^{\mathbb{B}} \upharpoonright A^{m(j)}$ for each $j \in J$.

Finally, an **automorphism** of an L -structure $\mathbb{A}$ is an isomorphism of $\mathbb{A}$ on itself; we write $\text{Aut}(\mathbb{A})$ for the group of automorphisms of $\mathbb{A}$.

5.2. Substructures. Let $X \subset B$ be a subset of the universe of some L -structure $\mathbb{B}$. The **substructure** of $\mathbb{B}$ **generated** by X is the smallest substructure of $\mathbb{B}$ containing X . Since the intersection of substructures is a substructure,

$$\mathbb{B}_X = \bigcap \{ \mathbb{A} \leq \mathbb{B} : X \text{ is a subset of the universe of } \mathbb{A} \}$$

We say that X **spans** $\mathbb{B}_X$. A structure $\mathbb{B}$ is **finitely generated** iff it is spanned by a finite subset. A structure $\mathbb{B}$ is **locally finite** iff all finitely generated substructures of $\mathbb{B}$ are finite.

Remark 5.1. Relational structures are locally finite.

5.3. Age of a structure. The **age** of a L -structure $\mathbb{A}$ is the class $\text{Age}(\mathbb{A})$ of all finite L -structures which are isomorphic to some substructure of $\mathbb{A}$.

5.4. Ultrahomogeneous structures. A locally finite structure $\mathbb{F}$ is **ultra homogeneous** iff each isomorphism between any two finite substructures of $\mathbb{F}$ can be extended to some automorphism of $\mathbb{F}$. A **Fraïssé structure** is an (infinite) countable ultrahomogeneous structure.

The following results (Proposition 5.2 and Theorems 5.3, 5.4) provide a characterization (Theorem 5.4) of all classes of finite structure which are the age of a Fraïssé structure. This can be done by using the method of back&forth; see [2, 10] for more details.

Proposition 5.2. *A locally finite L -structure $\mathbb{A}$ is ultrahomogeneous iff the following holds: If $\mathbb{B}, \mathbb{C} \in \text{Age}(\mathbb{A})$ and $\mathbb{B} \leq \mathbb{C}$ then each embedding $\mathbb{B} \longrightarrow \mathbb{A}$ can be extended to an embedding $\mathbb{C} \longrightarrow \mathbb{A}$.* $\square$

Theorem 5.3 (Fraïssé). *Any two (infinite) countable ultrahomogeneous L -structures having the same age are isomorphic.* $\square$

Theorem 5.4. *A non empty class of finite L -structures $\mathcal{C}$ is the age of a Fraïssé structure iff it satisfies:*

- (1) $\mathcal{C}$ is closed under isomorphisms: If $\mathbb{A} \in \mathcal{C}$ and $\mathbb{A} \cong \mathbb{B}$ then $\mathbb{B} \in \mathcal{C}$.
- (2) $\mathcal{C}$ is hereditary: If $\mathbb{A} \in \mathcal{C}$ and $\mathbb{B} \leq \mathbb{A}$ then $\mathbb{B} \in \mathcal{C}$.
- (3) $\mathcal{C}$ contains structures with arbitrarily high finite cardinality.
- (4) Joint embedding property: If $\mathbb{A}, \mathbb{B} \in \mathcal{C}$ then there is $\mathbb{D} \in \mathcal{C}$ such that $\mathbb{A} \leq \mathbb{D}$ and $\mathbb{B} \leq \mathbb{D}$.
- (5) Amalgamation property: Given $\mathbb{A}, \mathbb{B}_1, \mathbb{B}_2 \in \mathcal{C}$ and embeddings $\mathbb{A} \xrightarrow{f_i} \mathbb{B}_i$, $i \in \{1, 2\}$, there is $\mathbb{D} \in \mathcal{C}$ and embeddings $\mathbb{B}_i \xrightarrow{g_i} \mathbb{D}$ such that $g_1 \circ f_1 = g_2 \circ f_2$.

In such case there is a unique (up to isomorphism, infinite) countable Fraïssé structure $\mathbb{F}$ such that $\text{Age}(\mathbb{F}) = \mathcal{C}$; this $\mathbb{F}$ is the **Fraïssé limit** of $\mathcal{C}$ and we write $\mathbb{F} = \text{FLim}(\mathcal{C})$. $\square$

5.5. Fraïssé classes. A **Fraïssé class** is a class of finite structures $\mathcal{C}$ satisfying the conditions (1)...(5) of Theorem 5.4 above.

5.6. Colorings of structures. Given L -structures $\mathbb{A}, \mathbb{B}, \mathbb{C}$ we write

- $\text{Emb}(\mathbb{A}, \mathbb{B})$ for the set of embeddings from $\mathbb{A}$ to $\mathbb{B}$ en $\mathbb{B}$ as a substructure;
- and
- $\begin{pmatrix} \mathbb{A} \\ \mathbb{B} \end{pmatrix}$ for the set of substructures of $\mathbb{B}$ which are isomorphic to $\mathbb{A}$.

If $\mathbb{A}, \mathbb{B} \leq \mathbb{C}$; we also write $\mathbb{C} \longrightarrow (\mathbb{B})_r^{\mathbb{A}}$ whenever, for each $r \in \mathbb{N}$ and each r -coloring

$$c : \begin{pmatrix} \mathbb{C} \\ \mathbb{A} \end{pmatrix} \longrightarrow r$$

of the set $\begin{pmatrix} \mathbb{C} \\ \mathbb{A} \end{pmatrix}$, there is an isomorphic copy $\mathbb{B}' \leq \mathbb{C}$ of $\mathbb{B}$ such that $\begin{pmatrix} \mathbb{B}' \\ \mathbb{A} \end{pmatrix}$ is monochromatic.

Example 5.5. There is an obvious surjective map $\text{Emb}(\mathbb{A}, \mathbb{B}) \longrightarrow \left(\begin{smallmatrix} \mathbb{A} \\ \mathbb{B} \end{smallmatrix} \right)$. If $\mathbb{A}$ is a **rigid structure** (it does not admit non trivial automorphisms) then $\text{Emb}(\mathbb{A}, \mathbb{B})$ and $\left(\begin{smallmatrix} \mathbb{A} \\ \mathbb{B} \end{smallmatrix} \right)$ coincide.

Remark 5.6. An **order structure** is a structure in the language $L = (\leq)$ of ordered sets. If $\mathbb{A}, \mathbb{B}$ are two order structures and $\mathbb{A}$ is finite, then $\text{Emb}(\mathbb{A}, \mathbb{B})$ and $\left(\begin{smallmatrix} \mathbb{A} \\ \mathbb{B} \end{smallmatrix} \right)$ coincide.

5.7. Ramsey classes of structures. A Fraïssé class $\mathcal{C}$ has the **Ramsey property** iff, for each finite structure $\mathbb{B} \in \mathcal{C}$ and each substructure $\mathbb{A} \leq \mathbb{B}$, there is a finite structure $\mathbb{C} \in \mathcal{C}$ such that

$$(\forall r \in \mathbb{N}) \quad \mathbb{C} \longrightarrow (\mathbb{B})_r^{\mathbb{A}}$$

See [5, 6, 8, 12, 13, 14, 15, 16] for details and examples.

Lemma 5.7. *The group $\text{Aut}(\mathbb{F})$ of an infinite countable structure $\mathbb{F}$ is a closed subgroup of the infinite countable symmetric group S_∞ .*

Remark 5.8. Hence $\text{Aut}(\mathbb{F})$, with the induced topology as a subspace of S_∞ , is a polish group. On the other hand, if $\mathbb{A} \leq \mathbb{F}$ is a finite substructure and $n = |A|$ is the cardinality of its universe, then the action of S_∞ on $\mathbb{N}^{[n]}$ induces, in a natural way, an action of $\text{Aut}(\mathbb{F})$ on $\left(\begin{smallmatrix} \mathbb{F} \\ \mathbb{A} \end{smallmatrix} \right)$.

Theorem 5.9. *Let $\mathbb{F}$ be a Fraïssé structure and $\mathcal{C} = \text{Age}(\mathbb{F})$. The polish group $\text{Aut}(\mathbb{F})$ is extremely amenable if and only if $\mathcal{C}$ has the Ramsey property and all the structures of $\mathcal{C}$ are rigid.*

6. THE INFINITE RANDOM POLYHEDRON

6.1. Ordered polyhedra as a Ramsey class. It is easy to see that each pair $(x, S_x) \in \mathcal{P} \cup \mathcal{AP}$ is a relational structure whose universe is x and in which S_x is a countable family of relations over x . The notions of substructure, homomorphism, etc are induced by our geometric interpretation of the Ramsey space $\mathcal{P}$ (see Section 3). Furthermore, each one of these structures is rigid by construction. In particular, as we showed in Theorem 4.1 the following holds:

Theorem 6.1. *The class $\mathcal{AP}$ is a rigid Ramsey class of finite structures.* □

In virtue of of Theorem 5.9, we obtain the following:

Corollary 6.2. *Let $\mathbb{P} = \text{FLim}(\mathcal{AP})$, the Fraïssé limit of $\mathcal{AP}$. Then, $\text{Aut}(\mathbb{P})$ with the Polish topology inherited from S_∞ is extremely amenable.* □

6.2. The infinite random polyhedron. Consider a countably infinite set ω . We are going to define a family $S_\omega \subseteq \omega^{[<\infty]}$, as follows:

6.2.1. Hold a coin. Even a biased one will be fine, as long as the probability of each side of the coin is not 0. Define a family $T_\omega \subseteq \omega^{[<\infty]}$ probabilistically in the following way: for every $u \in \omega^{[<\infty]}$ such that $|u| > 1$ flip the coin, and say that u is in T_ω if and only if you get heads.

6.2.2. Set

$$(16) \quad S_\omega := \{\{x\} : x \in \omega\} \cup \{v : (\exists u \in T_\omega) v \subseteq u\}$$

It is easy to prove that S_ω is hereditary and $\bigcup S_\omega = \omega$. Hence (ω, S_ω) is an infinite polyhedron. We say that (ω, S_ω) is an *infinite countable random polyhedron*.

Lemma 6.3. *With probability 1, each finite polyhedron can be embedded in any infinite countable random polyhedron.*

[Proof] Without loss of generality, let us assume that the probability of getting a head in one flip of the coin is $\frac{1}{2}$. Fix a random polyhedron (ω, S_ω) , as constructed

in §6.2. Since $\omega^{[<\infty]}$ is infinite countable, there is a bijection $\varphi : \omega^{[<\infty]} \longrightarrow \mathbb{N}$. Then S_ω is a random event in the probability space

$$\Omega = \left(2^{\omega^{[<\infty]}} \cong 2^{\mathbb{N}}; \mathbb{B}(2^{\mathbb{N}}), P \right)$$

which is just the Cantor space $2^{\mathbb{N}}$ with the σ -algebra $\mathbb{B}(2^{\mathbb{N}})$ of Borel subsets and the probability measure P on $2^{\mathbb{N}}$ induced by the coin-flipping. By our assumptions; given a finite sequence $a \in 2^n$ the probability that the n th initial segment of an infinite countable random sequence of flips $z \in 2^{\mathbb{N}}$ coincides with a is

$$P(z \upharpoonright n = a) = \frac{1}{2^n}$$

which is the measure of the basic open subset $[a]$ in the product topology of $2^{\mathbb{N}}$; see equation (2) in §1. Via the Cantor map, Ω is isomorphic to the unit interval $[0, 1]$ with the Lébesgue measure. The key argument is that the probability of a singleton in Ω coincides with the Lébesgue measure of a point in $[0, 1]$, so it vanishes.

The bijection φ sends the family of faces of our random polyhedron to some infinite sequence $x = \varphi(S_\omega) \in 2^{\mathbb{N}}$. In order to show the statement of Lemma 6.3, by Lemma 3.1, it is enough to see that any finite simplex can be embedded in (ω, S_ω) . This is equivalent to say that (ω, S_ω) has substructures which are isomorphic to simplexes of arbitrarily big finite dimension; thus for any positive integer m there is some $n \geq m$ such that T_ω has some element u with cardinality $|u| = n$. Let us suppose the contrary; then there is some integer $m > 0$ such that $u \in T_\omega \Rightarrow |u| \leq m$.

This implies that the sequence $x = \varphi(S_\omega)$ has finitely many 1's and corresponds, via the Cantor map, to a rational number in $[0, 1]$; which is a contradiction since the probability of $[0, 1] \cap \mathbb{Q}$ is 0. $\square$

Remark 6.4. Though they are not isomorphic as countable structures, the random polyhedron and the random graph have the same probability distribution.

Theorem 6.5. *Let $\mathbb{P} = \text{FLim}(\mathcal{AP})$, the Fraïssé limit of $\mathcal{AP}$. Then $\mathbb{P}$ is an infinite ordered polyhedron which is isomorphic to (ω, S_ω) , as a polyhedron, and to $\mathbb{Q}$, as an ordered set.*

[Proof] By Lemma 6.3 we get $\text{Age}((\omega, S_\omega)) = \text{Age}(\mathbb{P}) = \mathcal{AP}$. $\square$

Corollary 6.6. *All infinite countable random polyhedra are isomorphic as countable structures.*

[Proof] By Theorem 5.3. □

The above facts allow us to we call (ω, S_ω) the **infinite random polyhedron**.

ACKNOWLEDGMENTS

The authors would like to thank A. Villaveces, P. Zambrano and the Seminario de Geometría y Teoría de Modelos (Seminario GeoMod) from the Universidad Nacional de Colombia at Bogotá and the Pontificia Universidad Javeriana at Bogotá, for helpful sessions, meetings and conversations. José Mijares would like to also thank Jesús Fernando Novoa, Head of the Department of Mathematics of the Pontificia Universidad Javeriana, for all the support during the realization of this research.

REFERENCES

- [1] BREDON, G. *Topology and Geometry*. Graduate Texts in Mathematics Vol. **139** Springer-Verlag (1993).
- [2] CHANG, C. & KEISLER, H. *Model Theory*. Studies in Logic and the Foundations of Math. Vol. **73** Elsevier- Amsterdam (1993).
- [3] CARLSON T. J. , SIMPSON S. G. **Topological Ramsey theory**. In: Nešetřil, J., Rödl (eds.) *Mathematics of Ramsey Theory*, 172-183, Springer (1990).
- [4] ELLENTUCK, E. *A new proof that analytic sets are Ramsey*. J. Symbol. Logic **39**, 163-165 (1974).
- [5] GRAHAM, R.L.; LEEB, K.; & ROTSCCHILD, B.L. *Ramsey's theorem for a class of categories*. Adv. Math. **8**, 417-433 (1972).
- [6] GRAHAM, R.L., ROTSCCHILD, B.L. & SPENCER j., *Ramsey Theory*, Wiley and sons. 1980.
- [7] KECHRIS, A. *Classical descriptive set theory*. Graduate Texts in Mathematics Vol. **156** Springer-Verlag (1994).
- [8] KECHRIS A. S., PESTOV V. G. and TODORCEVIC S., *Fraïssé Limits, Ramsey Theory, and topological dynamics of automorphism groups*, GEOMETRIC AND FUNCTIONAL ANALYSIS, Volume 15, Number 1 (2005), 106-189.
- [9] MAC LANE, S. *Categories for the working mathematician*. Graduate Texts in Mathematics Vol. **5**. Springer-Verlag. (1971).
- [10] MARKER, D. *Model theory, an introduction*. Graduate Texts in Mathematics. Vol. **217** Springer-Verlag. (2002).
- [11] MIJARES J. G., *On Galvin's Lemma and Ramsey spaces*. Ann. Comb. **16**, 319-330 (2012).
- [12] NEŠETRIL J., *Metric spaces are Ramsey*. European Journal of Combinatorics, **28**-#1, 457-468 (2007).
- [13] NEŠETRIL J., *Ramsey theory*. R.L. Graham, M. Grötschel, L. Lovász (Eds.), *Handbook of Combinatorics*, Elsevier (1995), 1331-1403.
- [14] NEŠETRIL J., RÖDL V., *Combinatorial partitions of finite posets and lattice -Ramsey lattices*, Algebra Universalis, **19**, 106-119 (1984).
- [15] NEŠETRIL J., RÖDL V., *Mathematics of Ramsey Theory*. Springer; 1990.
- [16] PESTOV, V. *Dynamics of infinite-dimensional groups*, ULECT 40, AMS, 2006.
- [17] PESTOV, V. *Mm-spaces and group actions*., L'Enseign. Math. **48**, 209-236 (2002).
- [18] RAMSEY, F. P. *On a problem of formal logic*. Proc. London Math. Soc. Ser. **2**#30, 264-286 (1929).
- [19] TODORCEVIC, S. *Introduction to Ramsey spaces*. Princeton University Press, Princeton, New Jersey, 2010.

DEPARTAMENTO DE MATEMÁTICAS, PONTIFICIA UNIVERSIDAD JAVERIANA, COLOMBIA; AND ESCUELA DE MATEMÁTICAS, UNIVERSIDAD CENTRAL DE VENEZUELA.

E-mail address: `jmijares@javeriana.edu.co`

E-mail address: `jose.mijares@ciens.ucv.ve`

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD NACIONAL DE COLOMBIA, K30-CL45, EDIF. 404, OFIC. 315-404. BOGOTÁ 101001000 COLOMBIA. (+571)3165000 EXT 13166

E-mail address: `gipadilla@unal.edu.co`